

Electronic supplementary information

Pd(II) and Pt(II) pincer complexes of a benzothiazole-appended methylthioacetamide ligand: synthesis and *in vitro* cytotoxicity

Diana V. Aleksanyan,^{a*} Svetlana G. Churusova,^a Ekaterina Yu. Rybalkina,^b
Yulia V. Nelyubina,^a and Vladimir A. Kozlov^a

^a *Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences,
ul. Vavilova 28, Moscow, 119991 Russia*

^b *Blokhin National Medical Research Center of Oncology of the Ministry of Health of the
Russian Federation, Kashirskoe shosse 23, Moscow, 115478 Russia*

*corresponding author: aleksanyan.diana@ineos.ac.ru

Table of contents

Fig. S1. ¹ H NMR spectrum of ligand 1 (400.13 MHz, CDCl ₃)	S2
Fig. S2. ¹³ C{ ¹ H} NMR spectrum of ligand 1 (100.61 MHz, CDCl ₃)	S3
Fig. S3. ¹ H NMR spectrum of complex 2 (400.13 MHz, CDCl ₃)	S4
Fig. S4. ¹³ C{ ¹ H} NMR spectrum of complex 2 (100.61 MHz, CDCl ₃)	S5
Fig. S5. ¹ H NMR spectrum of complex 3 (400.13 MHz, CDCl ₃)	S6
Fig. S6. ¹³ C{ ¹ H} NMR spectrum of complex 3 (100.61 MHz, CDCl ₃)	S7




Fig. S1. ^1H NMR spectrum of ligand 1 (400.13 MHz, CDCl_3)




Fig. S2. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of ligand 1 (100.61 MHz, CDCl_3)




Fig. S3. ¹H NMR spectrum of complex 2 (400.13 MHz, CDCl₃)




Fig. S4. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of complex **2** (100.61 MHz, CDCl_3)




Fig. S5. ^1H NMR spectrum of complex **3** (400.13 MHz, CDCl_3)




Fig. S6. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum of complex 3 (100.61 MHz, CDCl_3)