

Electronic supplementary information

SYNTHESIS AND CHARACTERIZATION OF TRIS(ALKOXY) YTTRIUM AND SCANDIUM COMPLEXES COORDINATED BY SCHIFF-BASE LIGANDS

A. I. Babkin,^a A. A. Kissel,^a Yu. V. Nelyubina,^a and A. A. Trifonov*^{a,b}

^a Nesmeyanov Institute of Organoelement Compounds, Russian Academy of Sciences,
ul. Vavilova 28, str. 1, Moscow, 119334 Russia

^b Razuvayev Institute of Organometallic Chemistry, Russian Academy of Sciences,
ul. Tropinina 49, Nizhny Novgorod, 603950 Russia

Table of contents

Table S1. Crystal data and structure refinement parameters for complex **1**

Figure S1. ^1H NMR spectrum (400 MHz, benzene- d_6 , 293 K) of **1**

Figure S2. $^{13}\text{C}\{\text{H}\}$ NMR spectrum (100 MHz, benzene- d_6 , 293 K) of **1**

Figure S3. ^1H NMR spectrum (400 MHz, benzene- d_6 , 293 K) of **2**

Figure S4. $^{13}\text{C}\{\text{H}\}$ NMR spectrum (100 MHz, benzene- d_6 , 293 K) of **2**

Table S1. Crystal data and structure refinement parameters for complex **1**

	1
Empirical formula	C ₆₉ H ₉₀ N ₃ O ₆ Y
Formula Weight	1146.34 g/mol
T, K	120
Crystal System	Monoclinic
Space Group	P2 ₁ /n
Unit Cell Dimensions	a = 12.538(9) Å b = 18.583(13) Å c = 30.54(2) Å α = 90° β = 99.552(15)° γ = 90°
V, Å ³	7016(9)
Z	4
d _{calc} , Mg/m ³	1.085
μ, mm ⁻¹	0.878
F ₀₀₀	2448
Crystal Size, mm	0.25x0.15x0.03
Θ Range for Data Collection, °	1.673-24.997
Index Ranges	-14<=h<=14, -22<=k<=22, -36<=l<=36
Reflns Collected	82567
Independent Reflns	12309
Observed Reflns	5028
Completeness to θ, %	99.8
Parameters	737
GOF on F ²	0.958
Final R Indices (I>2σ(I))	R ₁ = 0.00828, wR ₂ = 0.1754
R Indices (all data)	R ₁ = 0.2040, wR ₂ = 0.2359
Largest Diff Peak and Hole, e/Å ³	0.576 and -0.589




Figure S1. ^1H NMR spectrum (400 MHz, benzene- d_6 , 293 K) of **1**




Figure S2. $^{13}\text{C}\{^1\text{H}\}$ NMR spectrum (100 MHz, benzene- d_6 , 293 K) of **1**

